

Applying the Social Identity Approach to Public Health. A Case Example of Theory in
Practice

Abstract

Public health interventions play a major role in protecting, enhancing and maintaining the health of a population. At present, there is a plethora of public health interventions that promote a range of health behaviours for improving the lives of people suffering from chronic disease. Psychological theories can help us to understand the mechanisms that underpin health promotion interventions. However, the application and operationalisation of theory is not always transparent when utilised in public health systems for improving chronic health outcomes. Through the lens of social identity theory, the current article aims to provide a commentary on the content and practical delivery of a health intervention used within a public health service in the UK. Specifically, we present a case example detailing the application of social identity theory within a community pulmonary rehabilitation service to improve the quality of life for patients diagnosed with chronic obstructive pulmonary disease. Furthermore, we provide some balanced reflections on the utility of the social identity approach to inform a public health intervention. Our article provides practical applications and recommendations for using a novel theoretical approach in a public health setting to help bridge the theory to practice nexus.

Keywords: Group Membership, Exercise, Rehabilitation, Health

Applying the Social Identity Approach to Public Health. A Case Example of Theory in
Practice

Public health interventions aim to promote or protect health or prevent ill health in communities or populations and are complex. Complex interventions require theoretical underpinning to understand the mechanisms responsible for achieving change (Craig, Dieppe, Macintyre, Michie, Nazareth, & Petticrew, 2008). However, theory-based change techniques used in public health interventions are not always transparent. The obscurity of theoretically-derived techniques and their practical application can undermine the utility of public health interventions, creating a theory to practice nexus. As an example, descriptions of group-based change techniques often omit information about features relating to content and delivery (Borek, Abraham, Smith, Greaves, & Tarrant, 2015). The current article, therefore, provides an insight into the content and delivery of a group-based social identity intervention used within a community pulmonary rehabilitation service in the UK, which was based on a collaborative partnership between academic researchers and healthcare practitioners.

The social identity approach comprises of two related theories, namely social identity theory (Tajfel & Turner, 1979) and self-categorisation theory (Turner, 1985). At its core, the social identity approach stipulates that the way a person thinks and feels about self is derived from their group memberships. Group memberships can provide individuals with a sense of who they are in terms of a defined group identity (i.e., 'we' and 'us' rather than 'I' and 'me'). According to the social identity approach, two salient social processes are responsible for developing a group identity: 1) *categorisation*: awareness of similarities that collectively connect members as a group, and 2) *identification*: positively valuing the importance of belonging to a group membership. The social identity approach is particularly useful for informing interventions where people have limited access to social groups and networks (Haslam, Jetten, & Haslam, 2012). In public health settings, many chronically ill people often

experience social isolation (Johnson, Campbell, Bowers & Nichol, 2007) and impaired social interactions due to stigmatisation and as such have limited opportunities to develop new group memberships. The social identity approach, therefore, can usefully inform public health interventions that undertake programmes in group contexts (Steffens, La Rue, Haslam, Walter, Cruwys, Munt et al., 2019). Based on the social identity approach, recent research has developed an intervention for chronic obstructive pulmonary disease (COPD) patients attending the group-based exercise component of pulmonary rehabilitation (Levy, Matata, Pilsworth, Mcgonigle, Wigelsworth, Jones, et al., 2018). Preliminary findings suggest the social identity intervention improved COPD patients' social experiences and compliance with pulmonary rehabilitation (Pilsworth, Mcgonigle, Wigelsworth, Jones, Pott, Bettany, et al., 2018). To provide a further insight into the intervention's application, what follows next is a nuanced account of the content and delivery of social identity principles in a pulmonary rehabilitation setting.

The aim of the social identity intervention was to train healthcare practitioners to create a new group identity for COPD patients attending an 8-week pulmonary rehabilitation programme (see table 1). In view that COPD patients did not know each other before embarking upon pulmonary rehabilitation, the first phase of the intervention involved healthcare practitioners leading a 1-hour group discussion to create an awareness of a COPD pulmonary rehabilitation group membership. This process of categorisation accentuated the similarities between COPD patients and their differences from non-COPD sufferers to build the foundations of a COPD pulmonary rehabilitation ingroup. Establishing the COPD ingroup or the sense of 'us' was facilitated by the use of three categorisation techniques; 1) *self-anchoring*: projecting self-characteristics and attributes on the ingroup to provide meaning to one's new COPD pulmonary rehabilitation group membership, 2) *intergroup differentiation*: facilitating the differences between established COPD in-group attributes from those who do

not suffer from COPD (out-group) to provide a sense of unity among the COPD pulmonary rehabilitation group membership, and 3) *self-stereotyping*: healthcare practitioners projecting professional knowledge of collective COPD in-group characteristics and attributes on individuals that constitute the group membership to provide a sense of belonging. At each rehabilitation session, COPD patients wore specially designed uniform t-shirts that depicted the individual's name and an in-group logo. We believed the use of t-shirts facilitated a tangible sense of in-group distinctiveness to help facilitate the categorisation processes.

[INSERT TABLE 1 HERE]

The second phase of the intervention attempted to help COPD patients positively identify as being a part of the COPD group membership. This process of identification was necessary for COPD patients to internalise the concept of being a member of a group with a common purpose. We used group goal setting to help COPD patients identify with their group and subsequently develop a collective sense of 'we'. To this end, the setting of group goals provided COPD patients with a collective sense of fate and purpose (e.g., cohesion), direction and meaning, (e.g., achievement), effort and persistence (e.g., motivation). Healthcare practitioners implemented goal setting across three stages.

Stage one involved the setting of performance and process goals prior to pulmonary rehabilitation commencing. The former refers to collective group exercise performance attainments (e.g., collective rowing meters) and the latter refers to implementation strategies to help achieve collective performance goals (e.g., prosocial supportive behaviours). At stage two, healthcare practitioners discussed the group's goal progression through a 1-hour group discussion midway through the pulmonary rehabilitation programme. This stage allowed for the adjustment of goals to ensure they remained realistic and achievable. Some patients, for example, were likely to experience exacerbations that did affect rehabilitation attendance, completion and dropout, therefore requiring the adjustment of group goals. In stage three,

healthcare practitioners evaluated the group's goal achievements towards the end of pulmonary rehabilitation. During a 1-hour group supportive discussion meeting, healthcare practitioners were able to constructively communicate appropriate evaluative feedback by attributing goal successes to the group's collective endeavours (i.e., in-group internal bias), and attributing goal shortcomings to factors outside the group's control (i.e., in-group external bias). Additionally, healthcare practitioners were able to reinforce collective goal achievements by communicating personal gains from being a part of the group such as confidence and enjoyment (i.e., positive reinforcement) and how the group has helped remove unpleasant individual experiences such as isolation and anxiety (i.e., negative reinforcement).

Our article provides a case example of social identity processes in action to support COPD patients derive a sense of self during pulmonary rehabilitation as a member of a group. To date, the application of social identity has mainly focused on naturalistic identities (e.g., gender, nationality, sports teams) where the development of social identity processes and group memberships are ubiquitous. However, in public health contexts, naturalistic identities are not always prevalent, as was the case for COPD patients engaging with pulmonary rehabilitation. At present, there is a nexus between the social identity approach and public health practice, particularly with respect to developing new health-related group memberships and the social identities that exist within them. Bridging the theory-practice nexus is important for utilising the social identity approach for creating new group memberships in public health interventions. "There is nothing as practical as a good theory" (Lewin, 1952, p.169), therefore we encourage the greater publication of material that informs the application of social identity processes in the public health domain.

A caveat in the development of new social identities is the ability of individuals to sustain their group membership. This is an important issue as public health interventions are

often not sustainable due to being of finite duration resulting from funding restrictions (Walugembe, Sibbald, Le Ber, & Kothari, 2019). In the present case example, COPD patients attended an 8-week pulmonary rehabilitation programme and as such it was not possible for all patients to maintain their new group membership beyond this period. Potentially, this can be damaging for chronic disease populations, such as COPD patients, who can be often isolated and therefore unable to draw upon a broader network of group memberships. It is essential, therefore, to consider group membership continuity when developing new social identities for public health interventions. One fruitful area to explore is the role of peer-led support groups as an adjunct to public health service programmes to maintain group memberships and the multiple social identities that underpin them. Peer support groups have the potential to initiate a shared sense of identification that can facilitate the formation of constructive supportive relationships for people living with lifelong health conditions.

Healthcare practitioners have responsibility for leading pulmonary rehabilitation and are in a unique position to influence the social interactions of COPD patients. As such, healthcare practitioners can play an important role in delivering social identity-based interventions for public health services. Healthcare practitioners, in our intervention, undertook training on the application of social identity principles and processes within a pulmonary rehabilitation setting. However, this would not have been possible without a collaborative working partnership, alongside the development of relevant and bespoke training materials. Social identity interventions will have a greater chance of acceptance in public health settings with the establishment of effective partnerships between healthcare practitioners and the academic community, and when the application of the social identity approach has clear fit and relevance within existing health service provision.

Generally, when applying interventions within a public health service there is always the potential to experience unique organisational challenges. In our case, we encountered an

issue whereby, for theoretical purposes, we wanted rehabilitation groups to have uniform structure and composition to foster patient categorisation and identification. Doing so, however, resulted in delaying the start dates for some patients enrolling on a pulmonary rehabilitation programme. The challenge for us, therefore, was to ensure that we obtained a homogenous cohort of COPD patients in the social identity intervention within an appropriate waiting timeframe, as determined by the health service provider. We recommend that social-identity based interventions in public health will benefit from ascertaining, as early as possible, the organisational factors external to the intervention that might influence how it operates.

To conclude, this article provides a nuanced insight towards the application of the social identity approach in a public health setting. We attempt to bridge the gap between social identity principles and practice and in doing so raise some issues about maintaining social identities and the challenges of applying this approach in a public health context. The social identity approach holds promise to innovate and inform the development and sustainability of new and existing group structures across a spectrum of public health interventions. To realise this potential, we encourage the publication of applied and contextualised insights on the practicality and usability of the social identity approach in a variety of public health settings.

References

- Borek, A. J., Abraham, C., Smith, J. R., Greaves, C. J., & Tarrant, M. (2015). A checklist to improve reporting of group-based behaviour-change interventions. *BMC Public Health, 15*(1), 963. doi: 10.1186/s12889-015-2300-6
- Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2008). Developing and evaluating complex interventions: the new Medical Research Council guidance. *British Medical Journal, 337*, 979-983. doi: 10.1136/bmj.a1655
- Haslam, C., Jetten, J., & Haslam, A. (2012). Advancing the social cure: Implications for theory, practice and policy. In J. Jetten, C. Haslam & S.A Haslam (Eds.), *The social cure: identity, health and wellbeing* (pp. 319-343). Hove: East Sussex: Psychology Press.
- Hoffmann, T., Glasziou, P., Boutron, I., Milne, R., Perera, R., Moher, D et al. (2014). Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ, 348*, g1687. doi.org/10.1136/bmj.g1687
- Johnson, J.L., Campbell, A.C., Bowers, M., & Nichol, A.M. (2007). Understanding the social consequences of chronic obstructive pulmonary disease: the effects of stigma and gender. *Proceedings of the American Thoracic Society, 4*, 680–682. doi.org/10.1513/pats.200706-084SD.
- Levy, A.R., Matata, B., Pilsworth, S., Mcgonigle, A., Wigelsworth, L., Jones, L et al. (2018). An intervention for pulmonary rehabilitators to develop a social identity for patients attending exercise pulmonary rehabilitation: A feasibility and pilot randomised control trial protocol. *Pilot and Feasibility Studies, 4*, 40. doi: 10.1186/s40814-018-0238-z
- Lewin, K. (1952). *Field theory in social science: Selected theoretical papers by Kurt Lewin* (D. Cartwright Ed.). London, UK: Tavistock.
- Pilsworth, S., Mcgonigle, A., Wigelsworth, L., Jones, L., Pott, N., Bettany, M., et al. (2018).

Group-based social identity intervention during pulmonary rehabilitation improves COPD patients experience and promotes compliance with exercise programme.

Thorax, 73(4), 156. doi.org/10.1136/thorax-2018-212555.257

Steffens, N.K., La Rue, C.J., Haslam, C., Walter, Z.C., Cruwys, T., Munt, K.A., et al. (2019). Social identification-building intervention to improve health: A systematic review and meta-analysis. *Health Psychology Review*. Advance online publication. 10.1080/17437199.2019.1669481

Tajfel, H., & Turner, J.C. (1979). An integrative theory of intergroup conflict. In W.G. Austin & S. Worchel (Eds.), *The social psychology of intergroup relations* (pp. 33-47). Monterey, CA: Brooks/Cole.

Turner, J.C. (1985). Social categorisation and the self-concept: A social cognitive theory of group behaviour. In E.J. Lawler (Ed.), *Advances in group processes* (Vol. 2, pp. 77-122), Greenwich, CT: JAI Press.

Walugembe, D.R., Sibbald, S., Le Ber, M.J., & Kothari, A. (2019). Sustainability of public health interventions: Where are the gaps? *Health Research Policy and Systems*, 17, 8. doi.org/10.1186/s12961-018-0405-y