

Cognitive, Affective, & Behavioral Neuroscience

Implicit visual sensitivity towards slim versus overweight bodies modulates motor resonance in the primary motor cortex: a tDCS study

Journal:	<i>Cognitive, Affective, and Behavioral Neuroscience</i>
Manuscript ID	CABN-RA-19-230.R2
Manuscript Type:	Research Article
Date Submitted by the Author:	09-Oct-2020
Complete List of Authors:	MAKRIS, STERGIOS; Edge Hill University Cazzato, Valentina; Liverpool John Moores University, Psychology
Keywords:	Action observation, Primary Motor Cortex, tDCS, Motor resonance, Anti-fat attitudes

SCHOLARONE™
Manuscripts

1
2
3 **Implicit visual sensitivity towards slim versus overweight bodies modulates motor**
4
5 **resonance in the primary motor cortex: a tDCS study**
6
7

8 Stergios Makris^{ab} & Valentina Cazzato^c
9

10
11
12 ^aDepartment of Psychology, Edge Hill University, Ormskirk, UK
13

14
15 ^bArts and Wellbeing Research Centre, Edge Hill University, Ormskirk, UK
16

17
18 ^cSchool of Psychology, Faculty of Health, Liverpool John Moores University, Liverpool, UK
19
20

21
22
23
24
25
26
27 Correspondence: Dr Stergios Makris, Department of Psychology, Edge Hill University,
28 Ormskirk, Lancashire, L39 4QP, Phone: +44 1695 584102, Email: makriss@edgehill.ac.uk or
29
30 Dr Valentina Cazzato, School of Psychology, Faculty of Health, Tom Reilly Building, Byrom
31 Street, Liverpool, L3 3AF, Phone: +44 151 904 6208, Email: v.cazzato@ljmu.ac.uk
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Motor resonance (MR) can be influenced by individual differences and similarity in the physical appearance between the actor and observer. Recently, we reported that action simulation is modulated by an implicit visual sensitivity towards normal-weight, as compared to overweight bodies. Furthermore, recent research has suggested the existence of an action observation network responsible for MR, with limited evidence whether the primary motor cortex (M1) is part of this. Here, we aimed at expanding our previous findings with regards to the role of an implicit normal-weight-body preference in the MR mechanism, while at the same time we tested the functional relevance of M1 to MR, by using a transcranial direct current stimulation (tDCS) protocol. Seventeen normal-weight and 17 overweight participants were asked to observe normal-weight or overweight actors reaching and grasping a light or heavy cube and then, at the end of each video-clip to indicate the correct cube weight. Before the task, all participants received 15 mins of sham or cathodal tDCS over the left M1. Measures of anti-fat attitudes were also collected. During sham tDCS, all participants were better in simulating the actions performed by normal-weight, as compared to overweight models. Surprisingly though, cathodal tDCS selectively improved the ability in the overweight group to simulate actions performed by the overweight models. This effect was not associated with scores of fat phobic attitudes or implicit anti-fat bias. Our findings are discussed in the context of relevance of M1 to MR and its social modulation by anti-fat attitudes.

Keywords: action observation; primary motor cortex; tDCS; motor resonance; anti-fat attitudes

Introduction

There is mounting research evidence indicating that the observation of one person performing an action can automatically activate a network of cortical brain regions in the observer associated with action execution. This phenomenon is known as “motor resonance” (MR) and it is considered a critical element of one’s ability to anticipate forthcoming actions and make predictions about their outcome (Fadiga et al, 1995; Aglioti et al, 2008; Urgesi et al, 2010). Furthermore, it has been recently shown that MR can be modulated by one’s experience with the observed action (Buccino et al, 2004; Abreu et al, 2012), whereas physical and psychological similarities between the actor and observer can also influence the level of action simulation and anticipatory representation of other’s actions (Avenanti et al, 2010; Obhi et al, 2014; Azevedo et al, 2014; Cazzato & Makris, 2019).

The ability to perceive and simulate others’ actions is deemed as critical in social interactions and it is reciprocally linked to the empathic ability to understand someone’s intentions and emotions (Gapinski et al, 2006). For example, recent studies on pain perception have indicated that both physical similarity and group membership can influence the empathic resonant neural responses to others’ pain (Avenanti et al, 2010; Azevedo et al, 2013). More recently, we have shown that MR can also be modulated by differences in the body weight between the actor and observer and that explicit negative attitudes towards overweight models can influence our ability to accurately simulate and predict the outcome of their actions (Cazzato & Makris, 2019). This is an interesting finding indicating that not only previous experience with an observed action, but also the existence of a negative stereotype (see also Cazzato, Makris & Urgesi, 2017), can influence the level of one’s ability to perceive others’ actions and

1
2
3 **intentions. However, there is still limited evidence on how explicit and implicit anti-fat**
4 **attitudes can influence the MR, as well as the underlying neural mechanism.**
5
6
7
8
9

10 With regards to the neural underpinnings of MR, a plethora of neurophysiological studies
11 have indicated the existence of an action observation network (AON) involving mainly
12 visual, parietal and premotor regions (Caspers et al, 2010; Avenanti et al, 2013; Tidoni et al,
13 2013; Paracampo et al, 2017). Moreover, the primary motor cortex (M1) has been classically
14 considered to implement a mirror mechanism in perceiving and simulating others' actions.
15 However, there is so far elusive evidence whether it is **essential** for MR (Avenanti et al,
16 2007; Naish et al, 2016; Valchev et al, 2017). **In a recent seminal study, Paracampo and**
17 **colleagues (2018) aimed at investigating the causal role of M1 in human action**
18 **prediction by means of transcranial direct current stimulation (tDCS). Their results**
19 **have indicated that tDCS perturbation of the primary motor cortex (M1) could**
20 **diminish the subjects' ability to make accurate predictions about the observed actions.**
21 **This very interesting finding provides, for the first time, causal evidence about the role**
22 **of M1 in the MR mechanism, under diverse methodological parameters (task, tDCS**
23 **polarity, intensity, and site-specific disruption). However, whether individual**
24 **differences, or the existence of anti-fat attitudes could be linked to a modulation of the**
25 **MR mechanism under M1 stimulation goes beyond the aims of Paracampo's study.**
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Taking into consideration the aforementioned literature, in the present study, we aimed to
50 first of all replicate and expand our previous behavioural finding (Cazzato & Makris, 2019)
51 that the existence of a weight stereotype can influence the way one perceives and simulates
52 others' actions. Furthermore, we investigated whether M1 plays a critical role in the
53 simulation of observed actions as part of an extended AON. **Indeed, if M1 is essential for**
54
55
56
57
58
59
60

1
2
3 **action prediction, based on previous findings, we expected that modulating its neural**
4 **functioning by means of cathodal (disruptive) tDCS, would also disrupt performance in**
5 **the behavioural task.** To this aim, we have implemented the same object-weight
6 discrimination task (WDT, see also Finisguerra et al, 2018), in which participants had to
7 indicate whether the observed object (a cube) was light or heavy, along with a crucial
8 manipulation of the weight similarity between the model-actor and the observer-subject. It
9 has been previously shown that bluffing intentions as coded by incongruent kinematics may
10 result in a critical modulation of the MR effect (Tidoni et al, 2013; Finisguerra et al, 2018)
11 and as such, our task also involved two different types of action (truthful versus fake) to gain
12 a better insight on the mechanisms underlying MR. **Finally, the task was performed in two**
13 **counterbalanced sessions carried out immediately after active (cathodal tDCS) or sham**
14 **tDCS over left M1.** In keeping with previous studies indicating that cathodal currents over
15 the region reduce M1 excitability (Nitsche et al, 2008; Nitsche & Paulus, 2011; Stagg, Antal,
16 &Nitsche, 2018; Paracampo et al, 2018), we applied a similar protocol for the active tDCS
17 condition (cathodal tDCS, 2mA stimulation intensity, 15mins duration).

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40 **Implicit (or automatic) and explicit (self-report) measures of anti-fat attitudes were**
41 **measured by means of a weight implicit association test (weight-IAT) and the Fat**
42 **Phobia Scale, respectively. The rationale behind our methodological choice resided in**
43 **the fact that there are several interesting advantages in investigating both explicit and**
44 **implicit measures. Particularly, a number of empirical evidence suggests that people**
45 **often refrain from explicit endorsements of negative attitudes and stereotypes toward**
46 **social groups (Crosby et al., 1980; Greenwald et al., 2009; Teachman and Brownwell,**
47 **2001), thus implicit measures counter some of the limitations of explicit ones, such as**
48 **response bias and demand characteristics (see Gapinski et al. 2006). Furthermore,**
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **research showed that implicit attitudes predict certain forms of behaviour (e.g., non-**
4 **verbal behaviour and spontaneous behaviour), but not others (see Dovidio et al., 2002).**
5
6
7 **Hence, assessing both implicit and explicit facets of weight stigma may be deemed**
8 **necessary to an in-depth understanding of the correlates of anti-fat bias. Furthermore,**
9 **addressing the unique relationship between people's ability to predict others' actions**
10 **with both implicit and explicit anti-fat attitudes may inform development of**
11 **interventions for reducing anti-fat bias and for improving the health and well-being of**
12 **individuals, who are overweight or obese.**
13
14
15
16
17
18
19
20
21
22
23

24 **In agreement with our previous findings (Cazzato & Makris, 2019), we expected that**
25 **performance in the WDT, as an implicit measure of MR, would vary as a function of the**
26 **weight stigma, further corroborated by measures of weight-IAT and Fat phobia scale.**
27 **More specifically, we expected that at baseline (sham tDCS) both groups of participants**
28 **would perform better in the task after observing the normal-weight models, as**
29 **compared to the overweight ones. Finally, following up from previous evidence with**
30 **regards to the role of the primary motor cortex in the MR mechanism, we expected that**
31 **compared to sham tDCS, cathodal tDCS over M1 would disrupt, as implicitly measured**
32 **by performance in the task, the functionality of M1 and thus, highlight the critical role**
33 **of the primary motor cortex in the simulation and prediction of observed actions.**
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Methods

Participants

A total of 34 subjects (17 normal-weight, 7 females; 17 overweight; 13 females) participated in the study (mean age = 22 years, SD = 3.4). Participants were recruited internally through the Psychology SONA participation scheme at Liverpool John Moores University.

Additionally, participants were recruited externally through poster advertisements situated in public locations, social media and through individuals known to the researchers. As an incentive, participants either received SONA points and/or £10 in Shopping Vouchers. All subjects were recruited based on their body mass index (BMI). According to the World Health Organisation BMI criteria (WHO, 2000) those with a BMI between 18.5 – 24.9 were classified as normal-weight (mean = 22.4, SD = 1.5), whereas subjects with a BMI above 25 were classified as overweight (mean = 31.3, SD = 3.9). Participants' BMI was obtained from measuring weight (Kg) and height (cm), by means of a digital scale (OMRON BF511 full body composition scale) and a stadiometer. All subjects were right-handed, as measured by means of a standard handedness inventory (Briggs & Nebes, 1975; mean = 702.9, SD = 220.5), they reported normal or corrected-to-normal visual acuity and they were naïve to the purposes of the study. After providing an overview of the experimental procedure, including technical information about tDCS, all subjects provided written informed concern for participation. Before the tDCS session all subjects completed a medical screening questionnaire checking for contraindications to the use of tDCS (Rossi et al, 2009). They were all in good health, with no history of neurological or psychiatric disorders and free of psychotropic or vasoactive medication. After the end of the tDCS session none of the participants complained of any discomfort or adverse effects during the whole procedure.

Moreover, at the end of the experiment, participants filled the following self-report questionnaire (1) the Fat Phobia Scale—short form (Bacon, Scheltema, & Robinson, 2001), to measure explicit fat prejudice and completed (2) The Weight-Implicit Association Test (weight-IAT, Greenwald, Nosek, & Banaji, 2003) to assess implicit anti-fat bias, at the end of which they were debriefed as to the purposes of the study. As expected, an independent sample t-test indicated that BMI was significantly higher in overweight than in normal-weight participants. **However, the two groups were matched for age, Fat Phobia and Weight-IAT scores (see Table 1).** Finally, all the experimental procedures were approved by the Liverpool John Moores University research ethics committee and were in accordance with the ethical standards of the Declaration of Helsinki (1964).

General procedure

In two separate days, participants completed the WDT after cathodal or sham tDCS (counterbalanced within subjects and groups of normal weight and overweight participants) delivered over the left M1. Each block lasted for about 20mins (tDCS stimulation + task duration). Finally, after the tDCS experiment, participants were required to provide information about their weight and height (for calculating BMI) and to complete the Weight-IAT and the Fat Phobia Scale.

Weight discrimination task, stimuli, and kinematic analysis

The stimuli for the weight discrimination task were the same as in our previous study (see Cazzato & Makris, 2019). They consisted of a series of video clips depicting a normal-weight or overweight male or female model performing a reaching, grasping, and lifting an object

1
2
3 action (see Fig. 1A). The objects were two metal cubes of the same dimension ($5 \times 5 \times 5$ cm)
4
5 and identical appearance, but different weights (“light” approximately 100 gr and “heavy”
6
7 approximately 800 gr). All videos were recorded from the posterior plane with a Canon HD
8
9 Camera and they were then processed by means of the Adobe Premier Software (Adobe
10
11 Systems Incorporated, San Jose, CA) in order to appear in black and white, thus controlling
12
13 for local changes in skin tone, and have the same duration of 1600 ms (split in 8 frames, 200
14
15 ms each). During the first part of the recording the models were informed about the correct
16
17 weight of the cube and they were instructed to perform a congruent to the object’s weight
18
19 reaching, grasping and lifting action (true action, TA). For the second part, the actors were
20
21 also informed about the correct weight of the cube, but this time they were instructed to
22
23 perform an incongruent action (i.e. to pretend that the cube was light for the heavy one and
24
25 vice versa for the light one; fake action, FA). Overall, there were 16 video clips following a 4
26
27 actors (2 normal-weight, 2 overweight) \times 2 types of action (TA, FA) \times 2 metal cubes (light,
28
29 heavy) design. A subsequent kinematic analysis on the recorded video clips revealed
30
31 significant differences in the way models handled the different objects for the fake and true
32
33 trials, but only for the grasping and lifting actions and not the reaching one (for more
34
35 information on the kinematic analysis procedure and results see Cazzato & Makris, 2019).
36
37 This is quite important as we were expecting our subjects to make their judgments about the
38
39 weight of the depicted object based on the perceived kinematics during the grasping and
40
41 lifting phases.
42
43
44
45
46
47
48
49
50

51 During the experiment, all subjects were seated in a dimly lighted testing room in front of a
52
53 19 inches LCD monitor (resolution 1027×768 pixels, screen refresh frequency at 60 Hz).
54
55 The whole experiment was created in and controlled by E-Prime 2.0 Professional Software
56
57 (Psychology Software Tools, Pittsburgh, PA). At the beginning of the experimental session
58
59
60

1
2
3 subjects were required to provide their demographic information, then followed by brief
4
5 instructions about the task. Participants were reminded to ask any questions about the task,
6
7 and they completed an 8-trial practice block before proceeding with the actual experiment.
8
9 Each trial started with a centrally located black fixation cross presented on a grey
10
11 background. After one second, the video clips depicting the models performing the true or
12
13 fake action appeared for 1600 ms at the centre of the screen subtending a visual angle of
14
15 approximately $12^\circ \times 10^\circ$, followed by a question asking the participants to indicate whether
16
17 the object they saw in the trial was “heavy” or “light”. All 16 video clips were presented in a
18
19 random order controlled by the E-Prime software and overall, there were 3 blocks with 32
20
21 trials each (16 video clips repeated twice).
22
23
24
25
26
27
28

29 -----Please insert Figure 1 somewhere here -----
30
31
32

33 *Transcranial Direct Current Stimulation (tDCS) parameters*

34
35
36
37

38 tDCS was delivered using a battery-operated constant direct current stimulator (The Magstim
39
40 Co. Ltd., Whitland, Carmarthenshire, UK). A pair of surface sponge electrodes (5×5 cm)
41
42 were soaked with a standard saline solution (NaCl .9%) and held in place by elastic rubber
43
44 bands. To target left M1, the active electrode was placed over the C3 electrode position of the
45
46 10-20 system and the reference electrode was placed on the forehead over the contralateral
47
48 orbit area (Nitsche & Paulus, 2011) (see Fig.1B). Cathodal-tDCS was delivered with a
49
50 constant current of 2 mA. Stimulation lasted for 15 min, not including 20 seconds of ramp-up
51
52 and ramp-down at the beginning and end of stimulation, respectively. For sham tDCS, the
53
54 electrodes were placed on the same locations, but the current was turned on for only 30
55
56 seconds at the beginning of the sham session and was then turned off in a ramp-shaped
57
58
59
60

1
2
3 fashion (fade in/out: 20 sec). This way participants experienced the sensations initially
4 associated with the onset of stimulation (mild local tingling), without inducing any effective
5 modulation of cortical excitability. **Cathodal and sham tDCS sessions were**
6
7
8 **counterbalanced (for half of the participants the first tDCS session was sham followed**
9
10 **by cathodal tDCS on another day, and vice versa for the other half) and neither the**
11
12 **subject nor the researcher delivering the stimulation were informed about the type of**
13
14 **tDCS (double-blinded). As this was a within-subjects design, meaning that participants**
15
16 **received both sham and cathodal tDCS at different experimental sessions, an interval of**
17
18 **at least 48 hours was allowed between the two active and sham stimulation sessions, in**
19
20 **order to avoid carryover effects and to guarantee a sufficient washout of the effects of**
21
22 **the previous session (e.g., Mancini et al., 2012, Bolognini et al., 2010, Bolognini et al.,**
23
24 **2011).**
25
26
27
28
29
30
31
32

33 *Weight-IAT and Fat Phobia Scale*

34
35
36
37

38 At the end of the tDCS sessions, all participants completed **an implicit, automatic measure**
39 **of weight bias, i.e. Weight-IAT and one self-report, explicit measure of weight bias, by**
40 **means of the Fat Phobia Scale.** For the Weight-IAT, participants were required to answer as
41 fast and accurately as possible after the onset of the stimuli (i.e., single words or images), by
42 pressing a left (E) or a right (I) key on a computer keyboard with the index finger of the left
43 hand and right hand, respectively. The IAT lasted approximately 8 minutes and was
44 administered in seven blocks, consisting of both congruent and incongruent conditions
45 (blocks 3, 4, 6, and 7) and familiarization blocks (blocks 1, 2, and 5) (Greenwald, Nosek, &
46 Banaji, 2003; Cazzato, Makris, & Urgesi, 2017; Cazzato & Makris, 2019). Before the first
47 presentation of the weight-IAT, participants were shown a list with all the words belonging to
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the two relevant categories and were asked to carefully read all the stimuli. In the first block
4
5 of the weight-IAT, 12 images of overweight and 12 images of normal-weight people were
6
7 presented and had to be classified as being either 'Fat' (left key) or 'Normal-weight' (right
8
9 key) (Cazzato, Siega, & Urgesi, 2012). Each of the 12 images of the two categories was
10
11 presented only once for a total of 24 trials. The second block also consisted of 24 trials in
12
13 which **negative (requiring a left-key response) and positive (requiring a right-key**
14
15 **response) words were presented. Some examples for negative words (belonging to 'Bad'**
16
17 **category) are 'Terrible, Agony, and Horrible'. Some examples for positive words**
18
19 **(belonging to 'Good' category) are 'Joy, Wonderful, and Happy'. In the third block (24**
20
21 trials practice) and in the fourth block (48 trials test), both overweight and normal-weight
22
23 bodies and **good (positive) and bad (negative) words** were randomly presented, and
24
25 participants were instructed to press the left key for bad-related words and images of
26
27 overweight people, and the right key for good-related words and images of normal-weight
28
29 people (congruent-stereotype condition). In the fifth block (24 trials), response key
30
31 assignments were reversed in relation to the categorization involving images of overweight
32
33 people (right-key) and images of normal-weight people (left-key). Finally, in the sixth block
34
35 (24 trials practice) and in the seventh block (48 trials test), both overweight and normal-
36
37 weight bodies **and positive and negative words** were randomly presented and participants
38
39 were required to press the left key for images of overweight people and positive words and
40
41 the right key for images of normal-weight people and negative words (incongruent stereotype
42
43 condition). Typically, participants are faster and more accurate in the congruent- than in the
44
45 incongruent-stereotype blocks, thus demonstrating an automatic association between
46
47 overweight and 'Bad' categories and normal-weight and 'Good' categories (Greenwald,
48
49 Nosek, & Banaji, 2003). Stimuli were randomly presented within each block. Each
50
51 word/image remained on the computer screen until the participant gave a correct response in
52
53
54
55
56
57
58
59
60

1
2
3 each trial. Indeed, if an error occurred in a trial, a red “X” appearing below the word stimulus
4
5 prompted the participant to correct the mistake by pressing the correct key. Following the
6
7 response, the next stimulus appeared after 500 ms, during which only the category labels
8
9 were visible on the screen. The response latency data for each participant were transformed
10
11 into D scores using the D-algorithm, as developed by Greenwald, Nosek, & Banaji, (2003).
12
13 Accordingly, a positive D score indicates that a participant responded more quickly when
14
15 categorizing positive adjectives with ‘Normal-weight’ and negative with ‘Fat’, than when
16
17 categorizing in the opposite manner (‘Normal-weight’ with negative and ‘Fat’ with positive).
18
19
20
21
22
23

24 Finally, all participants filled the Fat Phobia Scale-short form self-report questionnaire
25
26 (Bacon, Scheltema, & Robinson, 2001), which assessed explicit negative attitudes and
27
28 stereotyped perceptions of obese people. In this measure, 14 pairs of adjectives are used to
29
30 describe obese people (e.g., “lazy” vs “industrious”, “no will power” vs “has will power”),
31
32 and respondents are asked to indicate on a scale from 1 to 5 which adjective they feel best
33
34 describes their beliefs about obese people. A score of 2.5 indicates neutral attitudes about
35
36 obese persons, with scores more than 2.5 reflecting higher levels of fat phobia (more negative
37
38 attitudes) and lower scores indicating more positive attitudes.
39
40
41
42
43
44

45 *Data handling*

46
47
48
49 Behavioural performance obtained at the two-alternative- forced-choice WDT after sham and
50
51 cathodal tDCS sessions was analysed using the signal detection theory (SDT). Based on SDT,
52
53 we calculated d' as a measure of (perceptual) sensitivity and $\ln\beta$ as a measure for the response
54
55 bias. The percentage of correct responses (accuracy) was first calculated for each participant
56
57 in each experimental condition. **In keeping with our previous study (Cazzato & Makris,**
58
59
60

1
2
3 **2019), the SDT parameters (d' and $\ln\beta$ scores) were calculated, considering whether**
4 **subjects' responses were congruent or not to the real weight of the cube. This way,**
5 **“heavy-object” responses to heavy-object stimuli were considered as hits and “heavy-**
6 **object” responses to light-object stimuli as false alarms. D' and $\ln\beta$ scores data from the**
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

WDT were then entered into a mixed four-way $2 \times 2 \times 2 \times 2$ ANOVA with: 2 (tDCS stimulations: cathodal tDCS, sham tDCS) \times 2 (model's weight: normal-weight, overweight) \times 2 (type of action: true, fake) as within-subject factors and the subject's weight (normal-weight, overweight) as a between-subject factor. Finally, we calculated, for each condition, a measure of the change of measure of perceptual sensitivity d' scores as the ratio between the individual values after cathodal tDCS and the corresponding values in the sham tDCS condition [cathodal tDCS/sham tDCS]. The change indices were correlated, using Pearson correlations, with individual scores obtained at the Fat Phobia Scale and the Weight-IAT. All statistical analyses were performed with STATISTICA 8.0 (StatSoft Inc, Tulsa, Oklahoma). The source of all significant repeated-measure ANOVA interactions was analysed using the Duncan's post-hoc tests. Effect sizes were estimated using the partial eta square variable (η^2). All data are reported as Mean (M) and Standard Error of the Mean (s.e.m.). A significance threshold of $p < 0.05$ was set for all effects.

Results

Implicit and explicit anti-fat attitudes

Further to the Weight-IAT, one-sample t tests were used to compare the mean D-scores to zero (where zero refers to the absence of any response bias) for both groups. Both normal-weight and overweight participants showed a significant stereotypical anti-fat bias, indicating that they were more prone to associate overweight people to the bad-related category and normal-weight people to the good-related category than vice versa [normal-weight group: $t(16) = 7.89, p < 0.001$; overweight group: $t(16) = 8.7, p < 0.001$]. Interestingly, the two groups did not differ in the levels of stereotypical explicit fat phobia. One-sample t tests were also performed to compare the mean Fat phobia scale scores to 2.5 (where 2.5 refers to a moderate level of explicit phobia against obese people) for each group. In accordance with implicit anti-fat bias results, both normal-weight and overweight participants showed high level of explicit negative attitudes and stereotyped perceptions of obese people [normal-weight group: $t(16) = 13.46, p < 0.001$; overweight group: $t(16) = 8.24, p < 0.001$].

WDT performance after sham and cathodal tDCS

The $2 \times 2 \times 2 \times 2$ ANOVA on the mean d' scores revealed main effects of subject's weight [$F(1,32) = 8.041, p < 0.001, \eta p^2 = 0.201$], model's weight [$F(1,32) = 31.510, p < 0.001, \eta p^2 = 0.496$] and of action type [$F(1,32) = 58.598, p < 0.001, \eta p^2 = 0.647$], further corroborated by a significant 2-way interaction between model's weight and action type [$F(1,32) = 123.547, p < 0.001, \eta p^2 = 0.794$], as well as a significant 2-way interaction of model's weight \times subject's weight [$F(1,32) = 9.352, p < 0.001, \eta p^2 = 0.226$]. Most

1
2
3 importantly, the 4-way interaction of tDCS stimulations, model's weight, action type
4 and subject's weight was also significant [$F(1,32) = 5.655, p = 0.024, \eta p^2 = 0.150$].

7
8 For truthful actions trials, post-hoc comparisons showed that compared to
9 sham stimulation, cathodal tDCS over M1 yielded no effects while normal-weight
10 participants were observing either overweight (sham tDCS: 1.22 ± 0.34 vs. cathodal
11 tDCS: $1.13 \pm 0.28, p = 0.564$) or normal-weight models (sham tDCS: 2.24 ± 0.37 vs.
12 cathodal tDCS: $2.26 \pm 0.34, p = 0.902$) performing a true action.

13
14
15
16
17
18
19 On the opposite, cathodal tDCS over M1 improved perceptual sensitivity when
20 overweight participants observed overweight models performing true actions, (sham
21 tDCS: 0.82 ± 0.34 vs. cathodal tDCS: $1.45 \pm 0.28, p = 0.001$). This effect was not detected
22 when overweight participants were observing normal-weight models performing a true
23 action (sham tDCS: 2.25 ± 0.37 vs. cathodal tDCS: $2.45 \pm 0.34, p = 0.258$).

24
25
26
27
28
29
30 For fake actions trials, when compared to sham stimulation, cathodal tDCS over M1
31 did not affect normal-weight participants' sensitivity scores while they were observing
32 either overweight (sham tDCS: -0.18 ± 0.35 vs. cathodal tDCS: $0.11 \pm 0.32, p = 0.08$) or
33 normal-weight models (sham tDCS: -2.39 ± 0.36 vs. cathodal tDCS: $-2.41 \pm 0.33, p =$
34 0.902) performing a fake action.

35
36
37
38
39
40
41
42 On the opposite, cathodal tDCS over M1 decreased the perceptual sensitivity whilst
43 overweight participants were observing an overweight model performing fake actions
44 (sham tDCS: 0.82 ± 0.34 vs. cathodal tDCS: $1.45 \pm 0.28, p = 0.026$). This effect was not
45 evident when overweight participants were observing normal-weight models
46 performing a fake action (sham tDCS: -2.34 ± 0.36 vs. cathodal tDCS: $-2.50 \pm 0.33, p =$
47 0.379).

48
49 -----Please insert Figure 2 somewhere here -----
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

To sum up, cathodal tDCS over M1 improved the ability of overweight participants to simulate the true actions when performed by the overweight models, and thus to make accurate responses in the task. Furthermore, cathodal tDCS improved their action simulation when observing fake actions performed by overweight models, as they were more fooled by their deceptive kinematics and thus, they made more error responses in the task. Finally, for $\ln\beta$ scores the ANOVA revealed no significant main effects or interactions (all $F_s < 3.8$; $p > 0.05$), confirming that the aforementioned effects were not mediated by a change in response bias.

Finally, no significant correlations were found between the tDCS indices and implicit weight bias (weight-IAT) or explicit fat phobic attitudes (Fat Phobia scale) scores for any of the models or group of participants.

Discussion

In the present study, we investigated how inhibitory versus control non-invasive brain stimulation over the left primary motor cortex (M1) can modulate the simulation and understanding of observed familiar actions in terms of a weight discrimination task. More specifically, we applied cathodal (inhibitory) and sham (control) tDCS over the M1 area of normal-weight and overweight participants, prior to observing normal-weight and overweight models performing truthful or fake reaching and grasping actions and making a decision about the weight of the object. Furthermore, at the end of the WDT, the two groups were required to complete the Weight-IAT and the Fat Phobic scale to investigate associations between explicit and implicit weight stereotypes and the effects obtained on the WDT after cathodal tDCS stimulation.

1
2
3
4
5
6 **In keeping with our previous (behavioural) study (Cazzato & Makris, 2019), the**
7
8 **statistical analysis of the WDT has revealed that during sham tDCS, both normal-**
9
10 **weight and overweight participants performed better when they observed the normal-**
11
12 **weight models performing the actions, as compared to the overweight ones. Most**
13
14 **importantly, tDCS over M1 has, surprisingly, improved the way that our overweight**
15
16 **participants were able to perceive and simulate the actions performed by the overweight**
17
18 **models. No such finding was observed in the normal-weight group or for the normal-**
19
20 **weight models. More specifically, for true actions, cathodal tDCS over M1 resulted in**
21
22 **better performance in the WDT when overweight participants observed the actions**
23
24 **performed by the overweight models, indicating that the inhibition of M1 resulted in an**
25
26 **increase of the level at which the participants could simulate the observed actions and**
27
28 **thus, make accurate judgments about the weight of the object. The opposite pattern was**
29
30 **observed for fake actions; i.e. their performance in the task was worse as compared to**
31
32 **sham stimulation, thus indicating that they were better into simulating the deceptive**
33
34 **kinematics, being fooled and making an inaccurate response. Finally, even though both**
35
36 **groups of normal- and overweight participants displayed similar and higher level of**
37
38 **explicit fat phobia and negative attitudes towards obese people, we did not find**
39
40 **correlational evidence to suggest an association between explicit and implicit weight**
41
42 **stereotypes and the effects obtained on the WDT after active tDCS stimulation. Despite**
43
44 **these unexpected effects, the aforementioned findings provide some further evidence on**
45
46 **the role of M1 in action observation and MR.**
47
48
49
50
51
52
53
54
55

56 Earlier studies on the neural basis of the motor resonance mechanism have described the
57
58 existence of an action observation network (AON) responsible for the simulation of observed
59
60

1
2
3 action sequences and the prediction of their outcome (Urgesi et al, 2010; Springer et al,
4
5 2013). Initially, in those studies the AON comprised mainly of visual, parietal and premotor
6
7 areas, whereas M1 was not considered part of the network, as previous neurophysiological
8
9 and neuroimaging studies did not reveal a consistent activation of M1 during action
10
11 observation (Gazzola & Keysers, 2009; Caspers et al, 2010). More recently, however, a series
12
13 of neurophysiological studies has revealed a modulation of neuronal activity in M1 during
14
15 action observation, similar to that of action execution, thus leading researchers to propose
16
17 that M1 could be considered part of an extended AON (Vigneswaran et al, 2013; Naish et al,
18
19 2014; Valchev et al, 2015). In a more recent study, Paracampo et al. (2018) have shown that
20
21 cathodal tDCS over M1 impaired the participants' accuracy in an action prediction task, but
22
23 only when the observed actions were performed by humans as compared to non-humans **and**
24
25 **thus, they provided seminal causal evidence of the critical role of M1 in action**
26
27 **simulation and prediction.**
28
29
30
31
32
33
34

35 **In the present study, we applied a tDCS protocol similar to that described by**
36
37 **Paracampo et al (2018) and our findings are somewhat comparable to those reported in**
38
39 **their paper, in the sense that cathodal stimulation over M1, as compared to sham, has**
40
41 **indeed modulated the way that our overweight participants perceived and simulated the**
42
43 **viewed actions. Moreover, our WDT was similar to their action prediction task, in the**
44
45 **sense that in both studies subjects had to accurately perceive the kinematics of the**
46
47 **models in order to make accurate predictions for the weight of the object (present**
48
49 **study) or the size of the object (Paracampo et al, 2018). A critical difference here,**
50
51 **though was that our methodological manipulation allowed for the investigation of**
52
53 **whether the existence of negative attitudes towards overweight models can modulate**
54
55 **performance in the task as a function of the MR mechanism. Indeed, despite the lack of**
56
57
58
59
60

1
2
3 **correlational evidence, both groups strongly endorsed negative stereotypes against fat**
4 **people, indicating that anti-fat bias can modulate the level of simulating and**
5 **understanding familiar actions performed by normal-weight or overweight people and**
6 **that, this effect on its own may be altered by the type of stimulation we applied over M1.**
7
8 **Hence, Paracampo et al (2018) were the first ones to provide causative evidence on the**
9 **critical role of M1 in the MR mechanism; however, the findings of the present study**
10 **have expanded on these results, by providing some further insight on how top-down**
11 **influences, such as the existence of anti-fat attitudes, can affect the MR mechanism and**
12 **the involvement of M1 in understanding others' actions.**
13
14
15
16
17
18
19
20
21
22
23
24
25

26 One consideration here is that cathodal tDCS only in our overweight subjects modulated the
27 simulation of fake or true actions performed by the overweight models. No such modulation
28 was observed in our slim subjects or for actions performed by the normal-weight models.
29
30

31 One could argue that if M1 is part of the AON and thus, involved in action perception and
32 prediction, tDCS should modulate the measured effects in the task, irrespective of the
33 physical characteristics of the actors, similar to the study of Paracampo and colleagues
34 (2018). **This is indeed a surprising finding and one explanation we could provide for**
35 **that is the fact that in the current study a critical methodological manipulation was the**
36 **effect of the anti-fat attitudes in simulating the actions of overweight people. In line with**
37 **previous findings (Cazzato & Makris, 2019), we have shown that during sham tDCS,**
38 **both normal-weight and overweight subjects endorsing high levels of explicit fat phobia**
39 **measures are worse at detecting and simulating the action kinematics of overweight**
40 **actors, as opposed to normal-weight ones. Hence, it could be that the simulation of the**
41 **actions performed by the normal-weight models had reached ceiling levels that could**
42 **not be further modulated by the tDCS.**
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Moreover, an alternative, but not mutually exclusive explanation to this could reside in other individual differences (which were not accounted for in this study), for example in the emphatic responses of our participants to normal-weight, as compared to overweight individuals. With these regards, there is increasing research evidence of an association between the magnitude of cortical activity in the AON and the primary motor cortex and empathic responses to others (Avenanti et al, 2005; Gazzola et al, 2006; Zaki et al, 2009; Perry et al., 2010). Recently a study by Jospe et al (2020) has indicated that manipulating levels of motor excitability, by means of non-invasive brain stimulation, can modulate the level of understanding and simulating others' actions. More specifically, they have shown that tDCS interference over M1 has impaired or improved their subjects' ability to perceive and simulate hand gestures, depending on their levels of empathy. Even though in the present study we have not measured or controlled for levels of empathy, it would not be surprising if both our normal-weight and overweight participants, due to their increased levels of anti-fat attitudes, would show less empathic responses for the overweight models (see also Lewis & Hodges, 2011). Therefore, according to the aforementioned literature, it is possible that tDCS over M1 could have improved the empathic resonance of our overweight subjects towards the actions of the overweight models (in group), thus improving their ability to simulate their action kinematics. Nevertheless, we argue that this is an assumption we have not tested here, and further research is deemed as necessary to validate this hypothesis.

Furthermore, there is increasing research evidence showing that M1 possesses functional and reciprocal connections with the somatosensory cortex (Gazzola &

1
2
3 **Keyzers, 2009; Bonini, 2017) and that SI is involved in perceiving proprioceptive**
4 **information from observed actions (Keyzers et al, 2004; Bolognini et al, 2011; Holle et**
5 **al, 2013). More recently, Valchev et al (2017) have shown that offline cTBS over SI, but**
6 **not M1, impaired the subjects' ability to accurately detect the weight of an object lifted**
7 **by an actor. They concluded that while SI is mainly involved in action perception by**
8 **extracting proprioceptive/tactile information derived from observed action kinematics,**
9 **the role of M1 in action simulation is rather debatable and many consider its activity to**
10 **be a simple downstream consequence of the reciprocal cortico-cortical connections with**
11 **SI (Geyer et al, 2000; Rizzolatti & Luppino; 2001). If this is the case, we hypothesize**
12 **that cathodal tDCS disruption of M1 may have reduced any noise in the network that**
13 **allowed our overweight participants to better simulate the observed actions from those**
14 **that they share the same or similar proprioceptive input, i.e. the overweight models. We**
15 **assume that the same effect was not observed in the normal-weight subjects, as their**
16 **performance in the task had already reached ceiling effects, irrespective of the type of**
17 **stimulation. Moreover, Valchev et al (2017) reported that disruption of M1 had variable**
18 **effects across their participants, as in half of their subjects it increased their**
19 **performance to the task. A similar finding has been also reported by Palmer et al.**
20 **(2016) where cTBS over M1 had variable effects across their participants when they**
21 **were observing familiar actions, leading to inhibition of M1 excitability in some and**
22 **increases in others. Nevertheless, the aforementioned studies suggest that the**
23 **observation of both implied and real action sequences can modulate sensorimotor**
24 **integration (i.e. Concerto et al, 2015) and even though in the present study we have not**
25 **tested for that, we cannot rule out that in some extent our findings could be the outcome**
26 **of a contribution of brain areas functionally connected to M1 (Mineo et al, 2018).**
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Finally, it should be noted here that, our main statistical approach in the present study**
4 **for analysing the subjects' performance in the task, was a measure of sensitivity into**
5 **detecting the appropriate kinematics, as opposed to measures of response accuracy.**
6
7 **More specifically, instead of just measuring how well the subjects performed the task**
8 **(response accuracy), by applying the signal detection theory we investigated the level at**
9 **which they were able to perceive differences in the observed kinematics, which was**
10 **critical for making accurate judgments in the task, but also for revealing the**
11 **involvement of the MR mechanism. Thus, we speculate that inhibitory tDCS over M1**
12 **could have altered the level of sensitivity for detecting accurate (true actions) or**
13 **incongruent (fake actions) kinematics, but only for the models that the subjects trusted**
14 **less, i.e. the overweight ones. Indeed, there is some research literature showing that the**
15 **anti-fat bias can make overweight people appear less trustworthy (Ravina, 2007;**
16 **Vartanian et al, 2014) and that could affect the level at which people perceive and**
17 **simulate their actions. If our hypothesis is correct, then tDCS interference over M1**
18 **could have altered the way participants approached and perceived the actions**
19 **performed by the overweight models, thus the increased sensitivity we have detected in**
20 **our results.**
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Some limitations should be considered, though, when interpreting the findings from this**
46 **study. The first one has to do with methodological limitation due to the low spatial**
47 **resolution of tDCS, as well as the montage we have applied for the reference electrode.**
48 **Even though traditionally tDCS has less spatial resolution as compared to other brain**
49 **stimulation techniques (i.e. transcranial magnetic stimulation, TMS), there is increasing**
50 **research literature showing that tDCS stimulation over M1 can successfully interfere**
51 **with the area's involvement in action perception and execution (i.e. Nitsche & Paulus,**
52
53
54
55
56
57
58
59
60

1
2
3 **2000, 2001). Nevertheless, some recent studies have indicated that response to tDCS**
4 **stimulation can be quite variable between participants (Nitsche & Paulus, 2001;**
5 **Tremblay et al, 2013), with some of them reporting that only 60% of subjects are**
6 **experiencing the classic tDCS interference effect, thus supporting the need for a method**
7 **of individualizing tDCS dosage which uses electric-field (E-field) modelling (Evans et al.**
8 **2020; Datta et al. 2012; Bikson, Rahman, and Datta, 2012), to allow for more consistent**
9 **responses to tDCS stimulation (Caulfield et al. 2020). We acknowledge that in the**
10 **present study we did not test our participants for a canonical response to M1 tDCS**
11 **prior to enrolling them and we agree that future tDCS studies should cautiously account**
12 **for this limitation. With regards to the montage of the reference electrode over the right**
13 **supraorbital area, it is thought that extracephalic electrode montages allow more focal**
14 **stimulation (Cogiamanian et al. 2007) and so that the location of the reference electrode**
15 **may have induced a spread of the cathodal current in anterior parts of the brain that**
16 **could have contaminated our findings (see Im et al. and 2012; Datta et al. 2009).**
17 **However, whilst a previous study from Avenanti et al. (2018) provided the first causal**
18 **evidence that the Inferior Frontal Cortex is involved not only in planning the execution**
19 **of an upcoming action, but also in making predictions about the outcomes of observed**
20 **actions, to the best of our knowledge, most indications point to the involvement of motor**
21 **and premotor cortexes in the understanding of other people actions (see for e.g., Zaki et**
22 **al., 2009; Avenanti et al., 2005; Gazzola et al., 2006) and not to the anterior prefrontal**
23 **cortex, whereas the fact that the current return from the electrode to the cortex**
24 **dispersed over a large area, would make it less likely to produce significant effects (see**
25 **also Jospe et al, 2020).**
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 In conclusion, in the present study we addressed the question of whether the anti-fat bias can
4 affect the way that we perceive and simulate the actions of other people, in line with the
5 theory of motor resonance. Moreover, we investigated how inhibitory non-invasive brain
6 stimulation over the primary motor cortex can modulate these effects. We have shown that
7 inhibitory tDCS over M1 can modulate motor resonance effects, thus providing further
8 evidence on the role of M1 in action perception and prediction. However, further studies on
9 this topic are deemed as necessary to validate, clarify and further expand those findings.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Compliance with Ethical Standards:**
4

5
6 **Funding:** This research did not receive any specific grant from funding agencies in the public,
7 commercial, or not-for-profit sectors.
8

9
10
11 **Conflict of Interest:** The authors declared no potential conflicts of interest with respect to the
12 research, authorship, and/or publication of this article.
13
14

15
16
17 **Ethical approval:** All procedures performed in studies involving human participants were in
18 accordance with the ethical standards of the institutional and/or national research committee
19 and with the 1964 Helsinki declaration and its later amendments or comparable ethical
20 standards. This article does not contain any studies with animals performed by any of the
21 authors.
22
23
24
25
26
27

28
29 **Informed consent:** Informed consent was obtained from all individual participants included
30 in the study.
31
32

33
34 **Acknowledgments:** We are grateful to Mr James Randle, Mr Alberto Failla and Ms Ashleigh
35 Bellard for helping with the data collection.
36
37
38

39
40 **Open practices:** all data and material are available upon reasonable request from the
41 corresponding authors.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References:

Abreu, A. M., Macaluso, E., Azevedo, R. T., Cesari, P., Urgesi, C., & Aglioti, S. M. (2012). Action anticipation beyond the action observation network: A functional magnetic resonance imaging study in expert basketball players. *European Journal of Neuroscience*, 35(10), 1646–1654.

Aglioti, S. M., Cesari, P., Romani, M., & Urgesi, C. (2008). Action anticipation and motor resonance in elite basketball players. *Nature Neuroscience*, 11(9), 1109–1116.

Avenanti, A., Buetti, D., Galati, G., & Aglioti, S. M. (2005). Transcranial magnetic stimulation highlights the sensorimotor side of empathy for pain. *Nature neuroscience*, 8(7), 955-960.

Avenanti, A., Bolognini, N., Maravita, A., & Aglioti, S. M. (2007). Somatic and motor components of action simulation. *Current Biology*, 17, 2129-2135.

Avenanti, A., Sirigu, A., & Aglioti, S. M. (2010). Racial bias reduces empathic sensorimotor resonance with other-race pain. *Current Biology*, 20(11), 1018–1022.

Avenanti A, Candidi M, Urgesi C. (2013). Vicarious motor activation during action perception: beyond correlational evidence. *Front Hum Neurosci*. 7.

1
2
3 **Avenanti, A., Paracampo, R., Annella, L., Tidoni, E., & Aglioti, S. M. (2018). Boosting**
4 **and decreasing action prediction abilities through excitatory and inhibitory tDCS of**
5 **inferior frontal cortex. *Cerebral Cortex*, 28(4), 1282-1296.**
6
7
8

9
10
11
12 Azevedo, R. T., Macaluso, E., Avenanti, A., Santangelo, V., Cazzato, V., & Aglioti, S. M.
13 (2013). Their pain is not our pain: Brain and autonomic correlates of empathic resonance
14 with the pain of same and different race individuals. *Human Brain Mapping*, 34(12), 3168–
15 3181.
16
17
18
19

20
21
22
23 Azevedo, R. T., Macaluso, E., Viola, V., Sani, G., & Aglioti, S. M. (2014). Weighing the
24 stigma of weight: An fMRI study of neural reactivity to the pain of obese individuals.
25 *Neuroimage*, 91, 109–119.
26
27
28
29

30
31
32
33 Bacon, J. G., Scheltema, K. E., & Robinson, B. E. (2001). Fat phobia scale revisited: The
34 short form. *International journal of obesity*, 25(2), 252.
35
36
37
38

39
40 **Bikson, M., Rahman, A., & Datta, A. (2012). Computational models of transcranial**
41 **direct current stimulation. *Clinical EEG and neuroscience*, 43(3), 176-183.**
42
43
44

45
46
47 **Bolognini, N., Olgiati, E., Rossetti, A., & Maravita A. (2010). Enhancing multisensory**
48 **spatial orienting by brain polarization of the parietal cortex. *European Journal of***
49 **Neuroscience**, 31, 1800-1806.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Bolognini, N., Rossetti, A., Casati, C., Mancini, F., & Vallar, G. (2011).**

4
5 **Neuromodulation of multisensory perception: a tDCS study of the sound-induced flash**
6
7 **illusion. *Neuropsychologia*, 49, 231-237.**

8
9
10
11
12 **Bonini, L. (2017). The extended mirror neuron network: Anatomy, origin, and**
13
14 **functions. *The Neuroscientist*, 23(1), 56-67.**

15
16
17
18
19 Briggs, G. G., & Nebes, R. D. (1975). Patterns of hand preference in a student population.
20
21 *Cortex*, 11(3), 230–238.

22
23
24
25
26 Buccino, G., Lui, F., Canessa, N., Patteri, I., Lagravinese, G., Benuzzi, F., et al. (2004).
27
28 Neural circuits involved in the recognition of actions performed by nonconpecifics: An
29
30 fMRI study. *Journal of Cognitive Neuroscience*, 16(1), 114–126.

31
32
33
34
35 Caspers, S., Zilles, K., Laird, A. R., & Eickhoff, S. B. (2010). ALE meta-analysis of action
36
37 observation and imitation in the human brain. *NeuroImage*, 50, 1148-1167.

38
39
40
41
42 **Caulfield, K. A., Badran, B. W., DeVries, W. H., Summers, P. M., Kofmehl, E., Li, X., ...**
43
44 **& George, M. S. (2020). Transcranial electrical stimulation motor threshold can**
45
46 **estimate individualized tDCS dosage from reverse-calculation electric-field modeling.**
47
48 ***Brain Stimulation*, 13(4), 961-969.**

49
50
51
52
53 Cazzato, V., Siega, S., & Urgesi, C. (2012). “What women like”: Influence of motion and
54
55 form on aesthetic body perception. *Frontiers in Psychology*, 3, 235.

1
2
3 Cazzato, V., Makris, S., & Urgesi, C. (2017). Cathodal transcranial direct current stimulation
4 of the extrastriate visual cortex modulates implicit anti-fat bias in male, but not female,
5
6 participants. *Neuroscience*, 359, 92–104.
7
8
9

10
11
12 Cazzato, V. & Makris, S. (2019). Implicit preference towards normal-weight bodies and
13 weight-stigma modulate the understanding of observed familiar actions. *Psychological*
14
15 *Research*, 83(8), 1825-1835.
16
17
18
19

20
21 **Cogiamanian, F., Marceglia, S. A. R. A., Ardolino, G., Barbieri, S., & Priori, A. (2007).**
22
23 **Improved isometric force endurance after transcranial direct current stimulation over**
24 **the human motor cortical areas. *European Journal of Neuroscience*, 26(1), 242-249.**
25
26
27
28
29

30
31 **Concerto, C., Al Sawah, M., Infortuna, C., Freedberg, D., Chusid, E., Aguglia, E., &**
32 **Battaglia, F. (2015). Neural circuits underlying motor facilitation during observation of**
33 **implied motion. *Somatosensory & motor research*, 32(4), 207-210.**
34
35
36
37
38
39

40 **Crosby, F., Bromley, S., & Saxe, L. (1980). Recent unobtrusive studies of Black and**
41 **White discrimination and prejudice: A literature review. *Psychological bulletin*, 87(3),**
42 **546.**
43
44
45
46
47
48

49 **Datta, A., Bansal, V., Diaz, J., Patel, J., Reato, D., & Bikson, M. (2009). Gyri-precise**
50 **head model of transcranial direct current stimulation: improved spatial focality using a**
51 **ring electrode versus conventional rectangular pad. *Brain stimulation*, 2(4), 201-207.**
52
53
54
55
56
57
58
59
60

1
2
3 **Datta, A. (2012). Inter-individual variation during transcranial direct current**
4 **stimulation and normalization of dose using MRI-derived computational models.**
5
6
7 **Frontiers in psychiatry, 3, 91.**

10
11
12 **Dovidio, J. F., Gaertner, S. E., Kawakami, K., & Hodson, G. (2002). Why can't we just**
13 **get along? Interpersonal biases and interracial distrust. Cultural Diversity and Ethnic**
14 **Minority Psychology, 8(2), 88.**

17
18
19
20
21 **Evans, C., Bachmann, C., Lee, J. S., Gregoriou, E., Ward, N., & Bestmann, S. (2020).**
22 **Dose-controlled tDCS reduces electric field intensity variability at a cortical target site.**
23 **Brain stimulation, 13(1), 125-136.**

24
25
26
27
28
29
30
31 Fadiga, L., Fogassi, L., Pavesi, G., & Rizzolatti, G. (1995). Motor facilitation during action
32 observation: A magnetic stimulation study. *Journal of Neurophysiology*, 73, 2608-2611.

33
34
35
36
37
38 Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power: A flexible statistical
39 power analysis program for the social, behavioral, and biomedical sciences. *Behavior*
40 *Research Methods*, 39, 175-191.

41
42
43
44
45
46
47 Finisguerra, A., Amoruso, L., Makris, S., & Urgesi, C. (2018). Dissociated representations of
48 deceptive intentions and kinematic adaptations in the observer's motor system. *Cerebral*
49 *Cortex*, 28(1), 33–47.

50
51
52
53
54
55
56 **Gapinski, K. D., Schwartz, M. B., & Brownell, K. D. (2006). Can television change anti-**
57 **fat attitudes and behavior? Journal of Applied Biobehavioral Research, 11(1), 1–28.**
58
59
60

1
2
3 **Gazzola, V., Aziz-Zadeh, L., & Keysers, C. (2006). Empathy and the somatotopic**
4 **auditory mirror system in humans. *Current biology*, 16(18), 1824-1829.**
5
6
7

8
9
10 Gazzola, V., & Keysers, C. (2009). The observation and execution of actions share motor and
11 somatosensory voxels in all tested subjects: Single-subject analyses of unsmoothed fMRI
12 data. *Cerebral Cortex*, 19, 1239-1255.
13
14
15
16

17
18
19 **Geyer, S., Schormann, T., Mohlberg, H., & Zilles, K. (2000). Areas 3a, 3b, and 1 of**
20 **human primary somatosensory cortex: 2. Spatial normalization to standard anatomical**
21 **space. *Neuroimage*, 11(6), 684-696.**
22
23
24
25

26
27
28 Greenwald, A. G., Nosek, B. A., & Banaji, M. R. (2003). Understanding and using the
29 implicit association test: I. An improved scoring algorithm. *Journal of Personality and Social*
30 *Psychology*, 85(2), 197.
31
32
33
34

35
36
37 **Greenwald, A. G., Poehlman, T. A., Uhlmann, E. L., & Banaji, M. R. (2009).**
38 **Understanding and using the Implicit Association Test: III. Meta-analysis of predictive**
39 **validity. *Journal of personality and social psychology*, 97(1), 17.**
40
41
42
43
44

45
46
47 Gutsell, J. N., & Inzlicht, M. (2010). Empathy constrained: Prejudice predicts reduced mental
48 simulation of actions during observation of outgroups. *Journal of Experimental Social*
49 *Psychology*, 46(5), 841–845.
50
51
52
53

1
2
3 Gutsell, J. N., Inzlicht, M. (2013). Using EEG mu-suppression to explore group biases in
4 motor resonance. In B. Derks, D. Scheepers, & N. Ellemers, (Ed.) Neuroscience of prejudice
5 and intergroup relations (pp. 279–297). Milton Park: Psychology Press.
6
7
8
9

10
11
12 **Holle, H., Banissy, M. J., & Ward, J. (2013). Functional and structural brain differences**
13 **associated with mirror-touch synaesthesia. *Neuroimage*, 83, 1041-1050.**
14
15
16
17

18
19 **Im, C. H., Park, J. H., Shim, M., Chang, W. H., & Kim, Y. H. (2012). Evaluation of local**
20 **electric fields generated by transcranial direct current stimulation with an extracephalic**
21 **reference electrode based on realistic 3D body modeling. *Physics in Medicine & Biology*,**
22 **57(8), 2137.**
23
24
25
26
27

28
29
30 **Jospe, K., Flöel, A., & Lavidor, M. (2020). The interactive effect of empathy and motor**
31 **cortex stimulation on hand gesture comprehension. *Neuropsychologia*, 107412.**
32
33
34
35

36
37 **Keysers, C., Wicker, B., Gazzola, V., Anton, J. L., Fogassi, L., & Gallese, V. (2004). A**
38 **touching sight: SII/PV activation during the observation and experience of touch.**
39 ***Neuron*, 42(2), 335-346.**
40
41
42
43
44

45
46
47 **Lewis, K. L., & Hodges, S. D. (2011). Empathy is not always as personal as you may**
48 **think: The use of stereotypes in empathic accuracy. *Empathy: From bench to bedside*,**
49 **73-85.**
50
51
52
53

54
55
56 **Mancini, F., Bolognini, N., Haggard, P., & Vallar, G. (2012). Tdcs modulation of**
57 **visually induced analgesia. *Journal of Cognitive Neuroscience*, 24, 2419-2427.**
58
59
60

1
2
3 **Mineo, L., Fetterman, A., Concerto, C., Warren, M., Infortuna, C., Freedberg, D., ... &**
4 **Battaglia, F. (2018). Motor facilitation during observation of implied motion: Evidence**
5 **for a role of the left dorsolateral prefrontal cortex. International Journal of**
6 **Psychophysiology, 128, 47-51.**
7
8
9
10
11
12
13

14 Naish, K. R., Houston-Price, C., Bremner, A. J., & Holmes, N. P. (2014). Effects of action
15 observation on corticospinal excitability: Muscle specificity, direction, and timing of the
16 mirror response. *Neuropsychologia*, 64, 331-348.
17
18
19
20
21

22
23
24 Naish, K. R., Barnes, B., & Obhi, S. S. (2016). Stimulation over primary motor cortex during
25 action observation impairs effector recognition. *Cognition*, 149, 84-94.
26
27
28
29

30
31 **Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human motor**
32 **cortex by weak transcranial direct current stimulation. The Journal of physiology,**
33 **527(3), 633-639.**
34
35
36
37
38
39

40 **Nitsche, M. A., & Paulus, W. (2001). Sustained excitability elevations induced by**
41 **transcranial DC motor cortex stimulation in humans. Neurology, 57(10), 1899-1901.**
42
43
44
45

46
47 Nitsche, M. A., Cohen, L. G., Wassermann, E. M., Priori, A., Lang, N., Antal, A., et al.
48 (2008). Transcranial direct current stimulation: State of the art 2008. *Brain Stimulation*, 1,
49 206-223.
50
51
52
53

54
55
56 Nitsche, M. A., & Paulus, W. (2011). Transcranial direct current stimulation-update 2011.
57 *Restorative Neurology and Neuroscience*, 29, 463-492.
58
59
60

1
2
3 Obhi, S. S., Hogeveen, J., Giacomin, M., & Jordan, C. H. (2014). Automatic imitation is
4 reduced in narcissists. *Journal of Experimental Psychology: Human Perception and*
5
6
7
8 Performance, 40(3), 920.
9

10
11
12 **Palmer, C. E., Bunday, K. L., Davare, M., & Kilner, J. M. (2016). A causal role for**
13
14 **primary motor cortex in perception of observed actions. *Journal of cognitive***
15
16 **neuroscience, 28(12), 2021-2029.**
17
18

19
20
21 Paracampo, R., Tidoni, E., Borgomaneri, S., di Pellegrino, G., & Avenanti, A. (2017).
22
23
24 Sensorimotor network crucial for inferring amusement from smiles. *Cerebral Cortex, 27,*
25
26 5116-5129.
27

28
29
30 Paracampo, R., Montemurro, M., de Vega, M., & Avenanti, A. (2018). Primary motor cortex
31
32 crucial for action prediction: A tDCS study. *Cortex, 109,* 287-302.
33
34

35
36
37 **Perry, A., Troje, N. F., & Bentin, S. (2010). Exploring motor system contributions to the**
38
39 **perception of social information: Evidence from EEG activity in the mu/alpha**
40
41 **frequency range. *Social neuroscience, 5(3), 272-284.***
42
43

44
45
46 **Ravina, E. (2007). Beauty, personal characteristics, and trust in credit markets.**
47
48 **Personal Characteristics, and Trust in Credit Markets (December 2007).**
49
50

51
52
53 **Rizzolatti, G., & Luppino, G. (2001). The cortical motor system. *Neuron, 31(6), 889-901.***
54
55
56
57
58
59
60

1
2
3 Rossi S, Hallett M, Rossini PM, Pascual-Leone A. 2009. Safety, ethical considerations, and
4 application guidelines for the use of transcranial magnetic stimulation in clinical practice and
5 research. *Clin Neurophysiol.* 120:2008–2039.
6
7
8
9

10
11
12 Springer, A., Parkinson, J., & Prinz, W. (2013). Action simulation: Time course and
13 representational mechanisms. *Frontiers in Psychology*, 4, 387.
14
15
16
17

18
19 Stagg, C. J., Antal, A., & Nitsche, M. A. (2018). Physiology of transcranial direct current
20 stimulation. *The Journal of ECT*, 34(3), 144-152.
21
22
23
24

25
26 **Teachman, B. A., & Brownell, K. D. (2001). Implicit anti-fat bias among health**
27 **professionals: is anyone immune?. *International journal of obesity*, 25(10), 1525-1531.**
28
29
30
31

32
33 Tidoni, E., Borgomaneri, S., Di Pellegrino, G., & Avenanti, A. (2013). Action simulation
34 plays a critical role in deceptive action recognition. *Journal of Neuroscience*, 33(2), 611–623.
35
36
37
38

39
40 **Tremblay, S., Beaulé, V., Lepage, J. F., & Théoret, H. (2013). Anodal transcranial direct**
41 **current stimulation modulates GABAB-related intracortical inhibition in the M1 of**
42 **healthy individuals. *Neuroreport*, 24(1), 46-50.**
43
44
45
46

47
48
49 Urgesi, C., Maieron, M., Avenanti, A., Tidoni, E., Fabbro, F., & Aglioti, S. M. (2010).
50 Simulating the future of actions in the human corticospinal system. *Cerebral Cortex*, 20(11),
51 2511–2521.
52
53
54
55
56
57
58
59
60

1
2
3 Valchev, N., Zijdwind, I., Keysers, C., Gazzola, V., Avenanti, A., & Maurits, N. M. (2015).
4
5 Weight dependent modulation of motor resonance induced by weight estimation during
6
7 observation of partially occluded lifting actions. *Neuropsychologia*, 66, 237-245.
8
9

10
11
12 Valchev, N., Tidoni, E., Hamilton, A. F., Gazzola, V., & Avenanti, A. (2017). Primary
13
14 somatosensory cortex necessary for the perception of weight from other people's action: A
15
16 continuous theta-burst TMS experiment. *NeuroImage*, 152, 195-206.
17
18

19
20
21 **Vartanian, L. R., Pinkus, R. T., & Smyth, J. M. (2014). The phenomenology of weight**
22
23 **stigma in everyday life. *Journal of Contextual Behavioral Science*, 3(3), 196-202.**
24
25

26
27
28 Vigneswaran, G., Philipp, R., Lemon, R. N., & Kraskov, A. (2013). M1 corticospinal mirror
29
30 neurons and their role in movement suppression during action observation. *Current Biology*,
31
32 23, 236-243.
33
34

35
36
37 World Health Organization. (2000). *Obesity: Preventing and managing the global epidemic*
38
39 (vol. 894). Geneva: World Health Organization.
40
41

42
43
44 **Zaki, J., Weber, J., Bolger, N., & Ochsner, K. (2009). The neural bases of empathic**
45
46 **accuracy. *Proceedings of the National Academy of Sciences*, 106(27), 11382-11387.**
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Captions:

Figure 1: (A) Sequence of presentation in a typical trial for the weight discrimination task (adapted from Cazzato & Makris, 2019). Pictures represent video-clip frames during which a) an overweight male model performs a true action, b) a normal weight male model performs a true action, c) an overweight female model performs a true action and d) a normal-weight female model performs a true action. (B) tDCS montage showing the positions of the active and reference electrodes on the left primary motor cortex (M1) and on contralateral orbit, respectively.

Figure 2: Mean (\pm SEM) of normal- and overweight models participants' task sensitivity (d'), during observation of true and fake actions performed by normal- and overweight-models upon the light or the heavy object during sham and cathodal tDCS (c-tDCS) over left M1. **A higher d' score corresponds to better task sensitivity.** Asterisks indicate significant comparisons ($p < 0.05$).

Table 1: Mean and Standard Deviation (SD, in brackets) of participants' demographics information and scores to the implicit and explicit measures of anti-fat attitudes (Weight-IAT and Fat Phobia Scale) as a function of groups' weight (Normal weight vs. Overweight).

	Normal weight	Overweight	Normal weight vs. Overweight
	n=17	n= 17	
Age	21.29 (3.58)	23.53 (2.92)	$t(32) = 1.99; p = 0.06$
BMI	22.43 (1.82)	31.34 (3.95)	$t(32) = 8.45; p < 0.001$
Weight-IAT	0.54 (0.28)	0.56 (0.26)	$t(32) = 0.17; p = 0.865$
Fat Phobia Scale	3.56 (0.33)	3.54 (0.52)	$t(32) = -0.16; p = 0.873$

Notes: The data of participants were compared by means of independent sample *t*-test; BMI, Body Mass Index; Weight-IAT, weight-implicit association test (D-score).





Figure 1: (A) Sequence of presentation in a typical trial for the weight discrimination task (adapted from Cazzato & Makris, 2019). Pictures represent video-clip frames during which a) an overweight male model performs a true action, b) a normal weight male model performs a true action, c) an overweight female model performs a true action and d) a normal-weight female model performs a true action. (B) tDCS montage showing the positions of the active and reference electrodes on the left primary motor cortex (M1) and on contralateral orbit, respectively.



Mean (\pm SEM) of normal- and overweight models participants' task sensitivity (d'), during observation of true and fake actions performed by normal- and overweight-models upon the light or the heavy object during sham and cathodal tDCS (c-tDCS) over left M1. A higher d' score corresponds to better task sensitivity. Asterisks indicate significant comparisons ($p < 0.05$).

317x247mm (96 x 96 DPI)